

5.4 The Binomial Probability Distribution

Definition. A **binomial experiment** is a random experiment that satisfies the following four conditions:

- There are n identical trials.
- The trials are independent, that is, the outcome of a trial doesn't affect the outcomes of the other trials.
- Each trial has two outcomes (often termed **success** and **failure**)
- The probability of success p is constant, that is, it doesn't change from trial to trial.

Example. Is flipping a fair coin 10 times a binomial experiment?

Answer. Yes.

- There is a predetermined number of flips ($n=10$).
- The flips are independent (assumed).
- Each trial results in either heads or tails.
- The probability of a success is $p=1/2$ (a fair coin) and is constant from trial to trial.

Example. The following are not binomial experiments:

- Flipping a fair coin until the first head appears (number of trials is not predetermined).
- Drawing one marble at a time without replacement from a box containing 8 white and 12 red marbles (probability of drawing a white, say, marble is not constant).
- Rolling a fair die 15 times (the number of outcomes is more than two).

Definition. Let a random variable X be the **number of successes** in a binomial experiment. Then X is called a **binomial random variable**, and it has a **binomial distribution** with parameters n and p , and the probability distribution

$$P(X = x) = {}_n C_x p^x (1 - p)^{n-x}$$
$$= \frac{n!}{x!(n-x)!} p^x (1 - p)^{n-x}, \quad x = 0, 1, \dots, n.$$

Proof. Consider a sequence of outcomes of n trials, of which x are successes and $n-x$ are failures. There are ${}_n C_x$ ways to find x places for the successes, and the other $n-x$ places will contain failures.

Each success has probability p and each failure has probability $1-p$, and the trials are independent. Therefore,

$$P(X = x) = {}_n C_x p^x (1-p)^{n-x}$$

A Note on Notation. The probability of failure is sometimes denoted by $q = 1 - p$, hence, the binomial probability may be written as

$$P(X = x) = {}_n C_x p^x q^{n-x}$$

Example. A fair coin is flipped 10 times. The flips are independent.

- Find the probability that exactly 6 heads appear.

Solution. Let X be the number of heads in the sequence of 10 flips. Then X has a binomial distribution with parameters $n=10$, and $p=0.5$. Therefore,

$$\begin{aligned} P(X = 6) &= {}_{10}C_6 (0.5)^6 (1-0.5)^{10-6} = \frac{10!}{6!(10-6)!} (0.5)^{10} \\ &= \frac{10 \cdot \overset{3}{\cancel{9}} \cdot \cancel{8} \cdot 7}{\cancel{4} \cdot \cancel{3} \cdot \cancel{2}} (0.5)^{10} = 210 (0.5)^{10} = 0.2051 \end{aligned}$$

- Find the probability that at most one head appears.

Solution.

$$\begin{aligned} P(X = 0) + P(X = 1) &= {}_{10}C_0 (0.5)^0 (0.5)^{10-0} + {}_{10}C_1 (0.5)^1 (0.5)^{10-1} \\ &= \frac{10!}{0!(10-0)!} (0.5)^{10} + \frac{10!}{1!(10-1)!} (0.5)^{10} \\ &= (1+10)(0.5)^{10} = 0.0107 \end{aligned}$$

Example. On a multiple-choice exam with three possible answers for each of the five questions, what is the probability that a student would get four or more correctly just by guessing? Let Y be the number of correct guesses on the exam. Then Y has a binomial distribution with parameters $n=5$, and $p=1/3$. We compute

$$\begin{aligned} P(Y \geq 4) &= P(Y = 4) + P(Y = 5) = {}_5 C_4 (1/3)^4 (2/3)^{5-4} \\ &+ {}_5 C_5 (1/3)^5 (2/3)^{5-5} = \frac{5!}{4!(5-4)!} (1/3)^4 (2/3) \\ &+ \frac{5!}{5!(5-5)!} (1/3)^5 = (5)(1/3)^4 (2/3) + (1/3)^5 = 0.0453 \end{aligned}$$

Definition. The **mean** of a **binomial random variable** X is $E(X) = np$.

Definition. The **variance** of a **binomial random variable** X is

$$\sigma^2 = np(1 - p) = npq.$$

Definition. The **standard deviation** of a **binomial random variable** X is

$$\sigma = \sqrt{np(1 - p)} = \sqrt{npq}.$$

Example. In our previous example,

- How many questions does the student expect to guess correctly?

Solution. $E(Y) = (5)(1/3) = 1.67.$

- What is the standard deviation of the number of correct guesses?

Solution.

$$\sigma = \sqrt{(5)(1/3)(1-1/3)} = \sqrt{10/9} = 1.0541.$$

Example. Screws produced by a company are defective with probability 0.2, independently of each other. A package of 12 screws is bought.

- What is the probability that at least one screw is defective?

Solution. Let X be the number of defective screws in the package. Then X is binomial with $n=12$, and $p=0.2$.

$$\begin{aligned} P(X \geq 1) &= 1 - P(X = 0) = 1 - {}_{12}C_0 (0.2)^0 (1-0.2)^{12} \\ &= 1 - (0.8)^{12} = 0.9313 \end{aligned}$$

- How many screws in the package we expect to be defective?

Solution. $E(X) = np = (12)(0.2) = 2.4$ defectives

- Find the standard deviation of the number of defectives in the package.

Solution.

$$\sigma = \sqrt{np(1-p)} = \sqrt{(12)(0.2)(1-0.2)} = 1.3856 \text{ defectives}$$

Example. A fair die is rolled four times.

- What is the probability that we obtain fewer than two 6's?

Solution. Let “rolling a 6” be a success, and let X be the number of successes in this experiment. Then X has a binomial distribution with $n=4$ and $p=1/6$. We compute

$$P(X < 2) = P(X = 0) + P(X = 1) = {}_4C_0(1/6)^0(1-1/6)^{4-0} \\ + {}_4C_1(1/6)^1(1-1/6)^{4-1} = (5/6)^4 + (4)(1/6)(5/6)^3 = 0.8681$$

- What is the expected number of 6's in the four rolls of a fair die?

Solution . $E(X) = np = (4)(1/6) = 0.67$ 6's

- What is the standard deviation of the number of 6's?

Solution. $\sigma = \sqrt{np(1-p)} = \sqrt{(4)(1/6)(1-1/6)} = 0.7454$ 6's

Example. A boy is lost in the woods. A search party is organized. Each member of the search party has probability 0.1 of finding the boy, independently of all others.

- How large should the search party be, so that the probability of finding the boy is at least 0.9999?

Solution. Let the search party be of size n .

Then $P(\text{finding the boy}) = 1 - P(\text{not finding the boy}) = 1 - (1 - 0.1)^n = 1 - (0.9)^n$.

For $n=87$, $1 - (0.9)^n = 0.999896$.

For $n=88$, $1 - (0.9)^n = 0.999906$.

Thus, the answer is that the search group should consist of a minimum of 88 people.

- Assuming the search party has 88 people, what is the average number of people who will find the boy?

Solution. $(88)(0.1) = 8.8$ people

- What is the standard deviation of the number of people who will find the boy?

Solution. $\sqrt{(88)(0.1)(1-0.1)} = 2.8$ people

Example. A bus-tour agency sold 12 tickets for a bus with 10 seats. Each person is a no-show with probability 0.2, independently of the others.

- Find the probability that all people who show up for the tour will be accommodated.

Solution. Let X be the number of people who show up. Then X has a binomial distribution with $n=12$ and $p=0.8$, and $P(\text{all are accommodated})=$

$$\begin{aligned} P(X \leq 10) &= 1 - P(X = 11) - P(X = 12) \\ &= 1 - (12)(0.8)^{11} (0.2) - (0.8)^{12} = 0.7251 . \end{aligned}$$

- How many people does the bus-tour agency expect will show up?

Solution. $E(X) = np = (12)(0.8) = 9.6$ people

- What is the standard deviation of the number of people who show up?

Solution.

$$\sigma = \sqrt{np(1-p)} = \sqrt{(12)(0.8)(1-0.8)} = 1.39 \text{ people}$$