Math 123: Taylor's Formula and Approximations

Ryan Blair

CSU Long Beach

Thursday October 31, 2013

Outline

Taylor's Formula

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + R_n(x)$$

Where $R_n(x)$ is the **error term of order n**.



$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + R_n(x)$$

Where $R_n(x)$ is the **error term of order n**.

Theorem (Taylor's Theorem)

Given a Taylor Series $\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k$, if there is a constant M such that $|f^{(n+1)}(t)| < M$ for all t between a and x, then $|R_n(x)| < M \frac{|x-a|^{n+1}}{(n+1)!}$

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + R_n(x)$$

Where $R_n(x)$ is the **error term of order n**.

Theorem (Taylor's Theorem)

Given a Taylor Series $\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k$, if there is a constant M such that $|f^{(n+1)}(t)| < M$ for all t between a and x, then $|R_n(x)| < M \frac{|x-a|^{n+1}}{(n+1)!}$

Uses: Can show Taylor series converges if $|R_n(x)|$ goes to zero as n goes to infinity, Can get estimates for functions.

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + R_n(x)$$

Where $R_n(x)$ is the **error term of order n**.

Theorem (Taylor's Theorem)

Given a Taylor Series $\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x-a)^k$, if there is a constant M such that $|f^{(n+1)}(t)| < M$ for all t between a and x, then $|R_n(x)| < M \frac{|x-a|^{n+1}}{(n+1)!}$

Uses: Can show Taylor series converges if $|R_n(x)|$ goes to zero as n goes to infinity, Can get estimates for functions.

Show that the Maclaurin series for cos(x) converges to cos(x) for all x using Taylor's Theorem.

Examples

- Show that the Maclaurin series for $\frac{1}{1-x}$ converges to $\frac{1}{1-x}$ for all $x \in [-\frac{1}{2}, \frac{1}{2}]$ by finding a formula for $R_n(x)$.
- ② Estimate the error for approximating e^x on $\left[\frac{-1}{2},\frac{1}{2}\right]$ using $1+x+\frac{x^2}{2}+\frac{x^3}{6}$.
- **3** Estimate the error for approximating cos(x) on $[-2\pi, 2\pi]$ using $1 + \frac{-x^2}{2} + \frac{x^4}{24}$.