

Math 123: Volumes and Arc Length

Ryan Blair

CSU Long Beach

Tuesday September 20, 2016

Outline

1 Volumes of Rotation

2 Arc Length

Volumes of solids of rotation

Replace all x 's with y 's in the following formulas to get other valid expressions for volume.

Disks:

$$\text{Vol} = \int_a^b \pi(\text{radius in terms of } x)^2 dx$$

Shells:

$$\text{Vol} = \int_a^b 2\pi(\text{radius in terms of } x)(\text{height in terms of } x) dx$$

Washers:

$$\text{Vol} = \int_a^b \pi(\text{outer radius in terms of } x)^2 - \pi(\text{inner radius in terms of } x)^2 dx$$

Volumes of solids of rotation

Replace all x 's with y 's in the following formulas to get other valid expressions for volume.

Disks:

$$\text{Vol} = \int_a^b \pi(\text{radius in terms of } x)^2 dx$$

Shells:

$$\text{Vol} = \int_a^b 2\pi(\text{radius in terms of } x)(\text{height in terms of } x) dx$$

Washers:

$$\text{Vol} = \int_a^b \pi(\text{outer radius in terms of } x)^2 - \pi(\text{inner radius in terms of } x)^2 dx$$

Exercise: Find the volume of the object obtained by rotating the region bounded by the curves $y = \cos(x) + 1$, $y = 0$ and $x = 0$ that contains $(1, 1)$ about the x -axis.

Volumes of solids of rotation

Replace all x 's with y 's in the following formulas to get other valid expressions for volume.

Disks:

$$\text{Vol} = \int_a^b \pi(\text{radius in terms of } x)^2 dx$$

Shells:

$$\text{Vol} = \int_a^b 2\pi(\text{radius in terms of } x)(\text{height in terms of } x) dx$$

Washers:

$$\text{Vol} = \int_a^b \pi(\text{outer radius in terms of } x)^2 - \pi(\text{inner radius in terms of } x)^2 dx$$

Exercise: Find the volume of the object obtained by rotating the region bounded by the curves $y = \cos(x) + 1$, $y = 0$ and $x = 0$ that contains $(1, 1)$ about the y -axis.

The length of a curve

Lets find the length of a curve by approximating by line segments.

The length of a curve

Lets find the length of a curve by approximating by line segments.

If f is continuous on the interval $[a, b]$, then the length of the graph of f from a to b is

$$L = \int_a^b \sqrt{1 + (f'(x))^2}$$

The length of a curve

Lets find the length of a curve by approximating by line segments.

If f is continuous on the interval $[a, b]$, then the length of the graph of f from a to b is

$$L = \int_a^b \sqrt{1 + (f'(x))^2}$$

Example: Find circumference of the circle $x^2 + y^2 = 4$.

The length of a curve

Lets find the length of a curve by approximating by line segments.

If f is continuous on the interval $[a, b]$, then the length of the graph of f from a to b is

$$L = \int_a^b \sqrt{1 + (f'(x))^2}$$

Example: Find circumference of the circle $x^2 + y^2 = 4$.

Example: Find the length of the curve $y = \ln(\cos(x))$ between $x = 0$ and $x = \frac{\pi}{3}$